


California Wildlife Habitat Relationships System

California Department of Fish and Game

California Interagency Wildlife Task Group

Chukar
Alectoris chukar
B132



Range maps are based on available occurrence data and professional knowledge. They represent current, but not historic or potential, range. Unless otherwise noted above, maps were originally published in Zeiner, D.C., W.F. Laudenslayer, Jr., K.E. Mayer, and M. White, eds. 1988-1990. California's Wildlife. Vol. I-III. California Depart. of Fish and Game, Sacramento, California. Updates are noted in maps that have been added or edited since original publication.

California Wildlife Habitat Relationships System
California Department of Fish and Game
California Interagency Wildlife Task Group

CHUKAR

Alectoris chukar

Family: PHASIANIDAE
B132

Order: GALLIFORMES

Class: AVES

Written by: G. Ahlborn
Reviewed by: N. Johnson
Edited by: G. Ahlborn

DISTRIBUTION, ABUNDANCE, AND SEASONALITY

Uncommon to common permanent resident of arid, rocky annual grassland and brush and scrub habitats with available water. First introduced into California in 1932. Established predominantly east of the Sierra Nevada and Cascade Range, and along inner Coast Ranges from Pacheco Pass to the Tehachapi Mts. Occurs in Great Basin and Mojave Desert vegetation associations. Found below sea level in Death Valley, and up to 3659 m (12,000 ft) in the White Mts. (Harper et al. 1958).

SPECIFIC HABITAT REQUIREMENTS

Feeding: Forages on the ground; gleans, grazes, and scratches. Primarily eats seeds of cheatgrass, other grasses, and exotic annual forbs. Also eats fruits and leaves of a variety of other plants. Adults consume less than 5% insects by volume; young eat more (Johnsgard 1973). Will dig through up to 20 cm (8 in) of snow for food.

Cover: Provided by rock outcrops, talus slopes, and vegetation. Roosts on slopes with at least 7% grade. Rocks and cliffs provide cover from inclement weather. Will fly or run, often upslope, to escape predators.

Reproduction: Nests often placed on steep talus slopes or slopes with rocky outcrops, above streams. Nest commonly placed under sagebrush, or in sagebrush-grassland, within 0.8 km (0.5 mi) of water.

Water: Congregates at water in summer. Drinks daily in hot, dry weather. Water is a major habitat component affecting distribution, especially in summer. In California, distribution generally limited to habitats that receive 12-50 cm (5-20 in) annual rainfall (Johnsgard 1973).

Pattern: Optimum habitat includes 25-50% steep, rocky or talus slopes, rock outcrops, cliffs and bluffs, and 50% sagebrush and cheatgrass, near brushy stream drainages with grasses and water (Galbreath and Moreland 1953, Christensen 1970).

SPECIES LIFE HISTORY

Activity Patterns: Yearlong, diurnal activity.

Seasonal Movements/Migration: Seasonal movements may include altitudinal migration of 8-16 km (5-10 mi) from lower elevations in winter to higher areas for summer.

Home Range: In Utah, coveys moved daily over 2.6 km² (1 mi²) (Phelps 1955). In dry periods, may travel 3.2 to 4.8 km (2-3 mi) to water (Johnsgard 1973).

Territory: Probably not territorial (Johnsgard 1973). In Washington, male defended only

area immediately around female during mating season (Mackie and Buechner 1963).

Reproduction: Breeds late March to early September; peak from mid-May to early July. Clutch size 10-20, average 15. Incubates about 24 days; hatching peaks in May. Male apparently assists with incubation and brooding; female may begin second clutch. Coveys of 5-40 form in late summer. Productivity varies greatly from year to year, affected mostly by weather.

Niche: Impact of chukar on native fauna and flora is unknown, although Harper et al. (1958) suggested that there was little competition for food with native species. Has been able to survive in habitats degraded by land use practices where native species have been extirpated locally. Cooper's hawks, coyotes, and ravens are among the numerous reported predators.

REFERENCES

- Christensen, G. C. 1970. The chukar partridge: Its introduction, life history, and management. Nevada Dep. Fish and Game, Reno. Biol. Bull. No. 4. 82pp.
- Ehrlich, P. R., D. S. Dobkin, and D. Wheye. 1988. The birder's handbook. Simon and Schuster, New York. 785pp.
- Galbreath, D. S., and R. Moreland. 1953. The chukar partridge in Washington. Wash. Dep. Game, Olympia. Biol. Bull. No. 11. 55pp.
- Grenfell, W. E., B. M. Browning, and W. E. Stienecker. 1980. Food habits of California upland game birds. Calif. Dep. Fish and Game, Sacramento. Wildl. Manage. Br. Admin. Rep. No. 80-1. 130pp.
- Harper, H. T., B. H. Harry, and W. D. Bailey. 1958. The chukar partridge in California. Calif. Fish and Game 44:5-50.
- Johnsgard, P. A. 1973. Grouse and quails of North America. Univ. Nebraska Press, Lincoln. 553pp.
- Leopold, A. S., R. J. Gutierrez, and M. T. Bronson. 1981. North American game birds and mammals. Scribner's, New York. 198pp.
- Mackie, R. J., and H. K. Buechner. 1963. The reproductive cycle of the chukar. J. Wildl. Manage. 27:246-260.
- Phelps, J. E. 1955. The adaptability of the turkish chukar partridge (*Alectoris graeca* Meisner) in central Utah. M.S. Thesis, Utah State Agric. Coll., Logan. 56pp. Dept. Inter. Bur. Land Manage., Riverside, CA (contact No. YA-512-CT8-4389)
- Remsen, J. V., Jr. 1978. Bird species of special concern in California. Calif. Dep. Fish and Game, Sacramento. Wildl. Manage. Admin. Rep. No. 78-1. 54pp.
- Skinner, M. P. 1938a. *Falco mexicanus* Schlegel: Prairie falcon. Pages 18-42 in A. C. Bent. Life histories of North American birds of prey. Part II. U.S. Natl. Mus. Bull. No. 170. 482pp.
- Smith, D. G., and J. R. Murphy. 1973. Breeding ecology of raptors in the eastern Great Basin of Utah. Brigham Young Univ., Provo. Sci. Bull. Biol. Ser. 18, No. 3. 76pp.
- Snow, C. 1974b. Prairie falcon, *Falco mexicanus*. U.S. Dep. Inter., Bur. Land Manage., Wash. DC. Habitat Manage. Ser. Rep. No. 8. 18pp.
- Steenhof, K. 1998. Prairie Falcon (*Falco mexicanus*) In, The Birds of North America, No. 346 (A. Poole and F. Gill, eds.). The Birds of North America, Inc., Philadelphia, PA.
- U.S. Bureau of Land Management. 1979. Snake River birds of prey special research report. U.S. Dep. Inter., Bur. Land Manage., Boise, ID. 142pp.
- Walton, B. J. 1977. Development of techniques for raptor management with emphasis on the peregrine falcon. Calif. Dep. Fish and Game, Sacramento. Admin. Rep. 77-4. 26pp.